

COSMOLOGICAL THEORIES OF SPECIAL AND GENERAL RELATIVITY - I

MOSHE CARMELI ^a

^a *Department of Physics, Ben Gurion University of the Negev, Beer Sheva 84105, Israel*

Abstract

In the standard cosmological theory one uses the Einstein concepts of space and time as were originally introduced for the special theory of relativity and the general relativity theory. According to this approach all physical quantities are described in terms of the continuum spatial coordinates and time. Using general relativity theory a great progress has been made in understanding the evolution of the Universe. Cosmologists usually measure spatial distances and redshifts of faraway galaxies as expressed by the Hubble expansion. In recent years this fact was undertaken to develop new theories in terms of distances and velocities (redshift). While in Einstein's relativity the propagation of light plays the major role, in the new theory it is the expansion of the Universe that takes that role and appears at the outset. The cosmic time becomes crucial in these recent theories, which in the standard theory is considered to be absolute but here it is relative. In this lecture this new approach to cosmology is presented.

1 Introduction

It is well known that both Einstein's theories are based on the fact that light propagates at a constant velocity. However, the Universe also expands at a constant rate when gravity is negligible. Moreover, cosmologists usually measure spatial distances and redshifts of faraway galaxies as expressed by the Hubble expansion. In recent years this fact was undertaken to develop new theories in terms of distances and velocities (redshift). While in Einstein's special relativity the propagation of light plays the major role, in the new theory it is the expansion of the Universe that takes that role. It is the concept of cosmic time that becomes crucial in these recent theories. In the standard theory the cosmic time is considered to be absolute. Thus we talk about the Big Bang time with respect to us here on Earth as an absolute quantity. Consider, for example, another galaxy that has, let us say, a relative cosmic time with respect to us of 1 billion year. Now one may ask what will be the Big Bang time with respect to this galaxy. Will it be the Big Bang time with respect to us minus 1 billion year? A person who lives in that galaxy will look at our galaxy and say that ours is far away from him by also 1 billion

year. Will that mean, with respect to him, our galaxy is closer to the Big Bang time by 1 billion year? Or will we seem to him to be farther by 1 billion year? All this leads to the conclusion that there is no absolute cosmic time. Rather, it is a relative concept.

Based on this assumption, we present in this lecture a theory that relates distances between galaxies and their relative velocities. These are actually the dynamical variables that astronomers measure. In the first part the theory will be relating distances to velocities with very weak gravitational field (special relativity). We then, in the second part, extend the theory to include the gravitational field of the Universe. Before doing that, we present a brief review of Einstein's special relativity theory.

2 Einstein's special relativity: A review

We will not give a full detail of this very important theory. Rather, we outline some of its fundamentals [1,2]

1. The Michelson-Morley experiment for the constancy of the speed of light.

2. Einstein's two postulates: (a) Constancy of the speed of light; (b) Validity of the laws of physics in internal coordinate systems. (See Figure 1)

3. The Lorentz transformation. This is given by

$$ct' = (ct - \beta x) / \sqrt{1 - \beta^2}, \quad x' = (x - \beta ct) / \sqrt{1 - \beta^2}, \quad (2.1)$$

where $\beta = v/c$, and $y' = y$, $z' = z$.

4. Minkowski's unification of space and time.

5. Invariance of the laws of physics under the Lorentz transformation.

6. Minkowski's line element

$$ds^2 = c^2 dt^2 - (dx^2 + dy^2 + dz^2) \quad (2.2)$$

7. The light cone (see Figure 2).

3 Cosmological special relativity

We outline this theory very briefly with the following points [3, 4, 5].

1. Hubble's law $R = \tau v$, where R is the distance to a galaxy, v is the receding velocity of the galaxy, τ is universal constant equal to 12.486 Gyr (Big Bang time).

2. Cosmic time is not absolute but a relative concept.

Example: Another galaxy is one billion year with respect to us. One may ask, what is the Big Bang time with respect to this galaxy. Will it be BB time with respect to us minus 1 billion year? From the point of view of that galaxy ours is far away from it by also 1 billion year. Does that mean, with respect to that galaxy, ours is closer to the BB time by 1 billion year? Or our galaxy will seem to be farther by 1 billion year? All this leads to the conclusion that there is no absolute cosmic time. Rather, it is a relative concept.




Figure 1: Two coordinate systems K and K' , one moving with respect to the other with a velocity v in the x -direction.

3. Universe with negligible gravity.

4. Line element of an expanding Universe with negligible gravity:

$$ds^2 = \tau^2 dv^2 - (dx^2 + dy^2 + dz^2). \quad (3.1)$$

It is equal to zero for the Hubble expansion but is not vanishing at cosmic times smaller than τ .

Notice the similarity to Minkowskian line element

$$ds^2 = c^2 dt^2 - (dx^2 + dy^2 + dz^2) \quad (3.2)$$

which vanishes for light propagation but is different from zero for particles of finite mass.

5. Postulates of cosmological special relativity: (1) The laws of physics are valid at all cosmic times; (2) τ has the same value at all cosmic times. (Similarly to Einstein's special relativity postulates.)

6. The cosmological transformation. This is the analog to the Lorentz transformation. It relates physical quantities at different cosmic times (similarly to the




Figure 2: The light cone in two dimensions, $x^0(= ct)$ and $x^1(= x)$. The propagation of two light signals in opposite directions passing through $x = 0$ at time $t = 0$, is represented by the two diagonal straight lines. (Compare the galaxy cone given in the sequel.)

Lorentz transformation that relates quantities at different velocities):

$$x' = \frac{x - (\tau - t)v}{\sqrt{(t/\tau)(2 - t/\tau)}}, \quad \tau v' = \frac{\tau v - x(\tau - t)/\tau}{\sqrt{(t/\tau)(2 - t/\tau)}}, \quad (3.3)$$

$$y' = y, \quad z' = z,$$

$0 \leq t \leq \tau$, $t = 0$ at the Big Bang, $t = \tau$, now.

7. Example: Denote the temperature of the Universe at a cosmic time t by T and that at present by T_0 ($=2.73\text{K}$), we then have

$$T = T_0 / \sqrt{(t/\tau)(2 - t/\tau)}. \quad (3.4)$$

At $t/\tau = 1/2$ we get $T = 3.15\text{K}$. (This result assumes negligible gravity and needs a correction by a factor of 13, and thus the temperature at $t/\tau = 1/2$ is 41K .)

8. The galaxy cone. This is the analog to the light cone in Einstein's special




Figure 3: The galaxy cone in cosmological relativity, describing the cone in the $x - v$ space satisfying $\tau^2 v^2 - x^2 = 0$, where x represents the three-dimensional space. The heavy dots describe galaxies. The galaxy cone represents the locations of the galaxies at a given time rather than their path of motion in the real space.

relativity. It represents the expansion of the Universe with negligible gravity. It is a four-dimensional cone. The coordinates are the velocity and the other three coordinates are the space coordinates. At the cone surface the Hubble expansion is represented, whereas the inner part of the cone represents events at cosmic times less than τ . The similarity to the light cone is remarkable.

9. Inflation at the early Universe. From the cosmological transformation we obtain the relationship between the mass density ρ_0 now to its value ρ at a backward time t :

$$\rho = \rho_0 / \sqrt{1 - t^2/\tau^2}. \quad (3.5)$$

The volume of the Universe is inversely proportional to its density, hence the ratio of the volumes at two backward cosmic times t_1 and t_2 is

$$V_2/V_1 = \sqrt{(1 - t_2^2/\tau^2) / (1 - t_1^2/\tau^2)}. \quad (3.6)$$

For t_1, t_2 very close to τ we have

$$V_2/V_1 = \sqrt{t'_2/t'_1}, \quad (3.7)$$

where primes indicate times with respect to the Big Bang. For $t'_2 - t'_1 \approx 10^{-32}\text{s}$, and t'_2 much less than 1s, we have

$$V_2/V_1 = 10^{-16}/\sqrt{t'_1}. \quad (3.8)$$

For example, at $t'_1 \approx 10^{-132}\text{s}$, we obtain $V_2/V_1 \approx 10^{50}$ describing inflation.

The above introduction gives a brief review of a new special relativity (cosmological special relativity, for more details see [5]). Obviously the Universe is filled up with gravity and therefore one has to go to a Riemannian space with the Einstein gravitational field equations in terms of space and redshift (velocity). This is done in the second part of these lectures. Before that we outline Einstein's general relativity theory.

4 General relativity theory: A brief outline

1. Postulates: (a) Principle of general covariance; (b) Principle of equivalence.
2. Riemannian curved space for the gravitational field. The metric tensor $g_{\mu\nu}$ as the gravitational potential [6].
3. The Einstein field equations:

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \kappa T_{\mu\nu} \quad (4.1)$$

4. The geodesic equation as the equation of motion:

$$\frac{d^2 x^\rho}{ds^2} + \Gamma_{\alpha\beta}^\rho \frac{dx^\alpha}{ds} \frac{dx^\beta}{ds} = 0. \quad (4.2)$$

5 Extension to curved space

The theory presented here, cosmological general relativity, uses a Riemannian four-dimensional presentation of gravitation in which the coordinates are those of Hubble, i.e. distances and velocity rather than the traditional space and time. We solve the field equations and show that there are three possibilities for the Universe to expand. The theory describes the Universe as having a three-phase evolution with a decelerating expansion, followed by a constant and an accelerating expansion, and it predicts that the Universe is now in the latter phase. It is shown, assuming $\Omega_m = 0.245$, that the time at which the Universe goes over from a decelerating to an accelerating expansion, i.e., the constant-expansion phase, occurs at 8.5 Gyr ago. Also, at that time the cosmic radiation temperature was 146K. Recent observations of distant supernovae imply that the Universe's growth is accelerating, contrary to what has always been assumed, that the expansion is

slowing down due to gravity. Our theory confirms these recent experimental results by showing that the Universe now is definitely in a stage of accelerating expansion. The theory predicts also that now there is a positive pressure, $p = 0.034g/cm^2$, in the Universe. It is worthwhile mentioning that the theory has no cosmological constant. It is also shown that the three-dimensional space of the Universe is Euclidean, as the Boomerang, Maxima, DASI and CBI microwave telescopes have shown. Comparison with general relativity theory is finally made and it is pointed out that the classical experiments as well as the gravitational radiation prediction follow from the present theory, too.

6 Cosmology in spacevelocity

In the framework of cosmological general relativity (CGR) gravitation is described by a curved four-dimensional Riemannian spacevelocity. CGR incorporates the BB constant τ at the outset. The Hubble law is assumed in CGR as a fundamental law. CGR, in essence, extends Hubble's law so as to incorporate gravitation in it; it is actually a *distribution theory* that relates distances and velocities between galaxies. The theory involves measured quantities and it takes a picture of the Universe as it is at any moment. The following is a brief review of CGR as was originally given by the author in 1996 [7].

The foundations of any gravitational theory are based on the principles of equivalence and general covariance. These two principles lead immediately to the realization that gravitation should be described by a four-dimensional curved spacetime, in our theory spacevelocity, and that the field equations and the equations of motion should be generally covariant. Hence these principles were adopted in CGR also. Use is made in a four-dimensional Riemannian manifold with a metric $g_{\mu\nu}$ and a line element $ds^2 = g_{\mu\nu}dx^\mu dx^\nu$. The difference from Einstein's general relativity is that our coordinates are: x^0 is a velocitylike coordinate (rather than a timelike coordinate), thus $x^0 = \tau v$ where τ is the Big Bang time and v the velocity. The coordinate $x^0 = \tau v$ is comparable to $x^0 = ct$ in ordinary general relativity. The other three coordinates x^k , $k = 1, 2, 3$, are spacelike, just as in general relativity theory.

An immediate consequence of the above choice of coordinates is that the null condition $ds = 0$ describes the expansion of the Universe in the curved spacevelocity (generalized Hubble's law with gravitation) as compared to the propagation of light in the curved spacetime in general relativity. This means one solves the field equations (to be given in the sequel) for the metric tensor, then from the null condition $ds = 0$ one obtains immediately the dependence of the relative distances between the galaxies on their relative velocities.

As usual in gravitational theories, one equates geometry to physics. The first is expressed by means of the Einstein tensor. The physical part is expressed by the energy-momentum tensor which now has a different physical meaning from that in Einstein's theory. More important, the coupling constant that relates geometry to physics is now also *different*.

Accordingly the field equations are

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \kappa T_{\mu\nu}, \quad (6.1)$$

exactly as in Einstein's theory, with κ given by $\kappa = 8\pi k/\tau^4$, (in general relativity it is given by $8\pi G/c^4$), where k is given by $k = G\tau^2/c^2$, with G being Newton's gravitational constant, and τ the Big Bang constant time. When the equations of motion are written in terms of velocity instead of time, the constant k will replace G . Using the above equations one then has $\kappa = 8\pi G/c^2\tau^2$.

The energy-momentum tensor $T^{\mu\nu}$ is constructed, along the lines of general relativity theory, with the speed of light being replaced by τ . If ρ is the average mass density of the Universe, then it will be assumed that $T^{\mu\nu} = \rho u^\mu u^\nu$, where $u^\mu = dx^\mu/ds$ is the four-velocity. In general relativity theory one takes $T_0^0 = \rho$. In Newtonian gravity one has the Poisson equation $\nabla^2\phi = 4\pi G\rho$. At points where $\rho = 0$ one solves the vacuum Einstein field equations in general relativity and the Laplace equation $\nabla^2\phi = 0$ in Newtonian gravity. In both theories a null (zero) solution is allowed as a trivial case. In cosmology, however, there exists no situation at which ρ can be zero because the Universe is filled with matter. In order to be able to have zero on the right-hand side of (6.1) one takes T_0^0 not as equal to ρ , but to $\rho_{eff} = \rho - \rho_c$, where ρ_c is the critical mass density, a *constant* in CGR given by $\rho_c = 3/8\pi G\tau^2$, whose value is $\rho_c \approx 10^{-29}g/cm^3$, a few hydrogen atoms per cubic meter. Accordingly one takes $T^{\mu\nu} = \rho_{eff}u^\mu u^\nu$; $\rho_{eff} = \rho - \rho_c$ for the energy-momentum tensor. Moreover, the above choice of the energy-momentum tensor is the only possibility that yields a constant expansion when $\rho = \rho_c$ as it should be.

In Part II we apply CGR to obtain the accelerating expanding Universe and related subjects.

References

- [1] A. Einstein, *Autobiographical Notes*, P.A. Schlipp, Ed., Open Court (1979)
- [2] A. Einstein, *The Meaning of Relativity*, 5th Edition, Princeton Univ. Press, Princeton (1955)
- [3] M. Carmeli, *Found. Phys.* **25**, 1029 (1995)
- [4] M. Carmeli, *Found. Phys.* **26**, 413 (1996)
- [5] M. Carmeli, *Cosmological Special Relativity*, 2nd Edition, World Scientific (2002)
- [6] M. Carmeli, *Classical Fields: General Relativity and Gauge Theory*, Wiley, New York (1982) [reprinted by World Scientific (2001)]
- [7] M. Carmeli, *Commun. Theor. Phys.* **5**, 159 (1996)